

**2024/TDC (CBCS)/EVEN/SEM/
MTMHCC-402T/233**

TDC (CBCS) Even Semester Exam., 2024

MATHEMATICS

(4th Semester)

Course No. : MTMHCC-402T

(Riemann Integration and Series of Functions)

Full Marks : 70

Pass Marks : 28

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

UNIT—I

1. Answer any two of the following questions :

2×2=4

- (a)** Write two partitions of the interval $[0, 1]$ such that one of them is a refinement of the other.
- (b)** Give example with justification of a bounded function on $[0, 2024]$ which is not Darboux integrable.

(2)

- (c) Define Riemann sum of a function with respect to a tagged partition of the domain $[a, b]$. Hence, define Riemann integral.

2. Answer either [(a) and (b)] or [(c) and (d)] : 10

- (a) Show that a bounded function f on $[a, b]$ is Darboux integrable iff for every $\varepsilon > 0$, there exists a partition P of $[a, b]$ such that $U(P, f) - L(P, f) < \varepsilon$. 5

- (b) Show that Riemann integral of a function, if it exists, is unique. 5

- (c) If P and P^* are two partitions of $[a, b]$ where P^* is a refinement of P , show that

$$U(P^*, f) - L(P^*, f) < U(P, f) - L(P, f) \quad 5$$

- (d) Show that every monotone function on $[a, b]$ is Riemann integrable. 5

UNIT—II

3. Answer any two of the following questions :

2×2=4

- (a) If $f(x) \geq 0 \forall x \in [a, b]$, show that

$$\int_a^b f dx \geq 0$$

(3)

- (b) If f is integrable on $[a, b]$, and g is not integrable on $[a, b]$, what can you conclude about the integrability of $f + g$ on $[a, b]$? Justify your answer.

- (c) Give example with justification, of a function f such that $|f|$ is integrable but f is not integrable.

4. Answer either [(a) and (b)] or [(c) and (d)] : 10

- (a) Show that the sum of two integrable functions is integrable. 5

- (b) Let f be bounded and integrable on $[a, b]$ and there exists a function F on $[a, b]$ such that $F' = f$ on $[a, b]$, then show that

$$\int_a^b f dx = F(b) - F(a) \quad 5$$

- (c) Show that if f is integrable on $[a, b]$, then it is also integrable on $[a, c]$ and $[c, b]$ where $c \in (a, b)$. Also show that

$$\int_a^b f dx = \int_a^c f dx + \int_c^b f dx \quad 5$$

- (d) If f and g are integrable on $[a, b]$ and g keeps the same sign over $[a, b]$, then prove that there exists a number μ lying between the bounds of f such that

$$\int_a^b fg dx = \mu \int_a^b g dx \quad 5$$

(4)

UNIT—III

5. Answer any *two* of the following questions :

2×2=4

(a) Test the convergence of

$$\int_0^1 \frac{dx}{\sqrt{x}}$$

(b) Write the comparison test for testing the convergence of $\int_a^b f dx$ at a , in limit form.

(c) Evaluate :

$$\int_0^1 x^4 (1-x)^5 dx$$

6. Answer *either* [(a) and (b)] or [(c) and (d)] : 10

(a) Examine the convergence of

$$\int_{-1}^1 \frac{dx}{(2-x)\sqrt{1-x^2}}$$

and evaluate the value, in case it converges.

5

(b) Show that

$$\int_0^1 x^{m-1} (1-x)^{n-1} dx$$

converges iff $m, n > 0$.

5

(5)

(c) Find the values of p for which

$$\int_0^{\pi/2} \frac{\sin x}{x^p} dx$$

converges.

4

(d) Prove that

$$\int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} dx = \beta(m, n)$$

and hence show that

$$\int_0^1 \frac{x^{m-1} + x^{n-1}}{(1+x)^{m+n}} dx = \beta(m, n)$$

6

UNIT—IV

7. Answer any *two* of the following questions :

2×2=4

(a) Define uniform convergence of a sequence of functions on the closed interval $[a, b]$. Illustrate with an example.(b) Check whether the sequence of functions $\langle x^n \rangle$ on $[0, 1]$ converges uniformly.(c) Use Weirstrass M -test to check the convergence of the series

$$\sum_{n=1}^{\infty} \frac{\sin nx}{n^3}$$

(6)

8. Answer either [(a) and (b)] or [(c) and (d)] : 10

(a) Show that the sequence of functions

$$\left\langle \frac{x}{x+n} \right\rangle$$

converges uniformly in $[0, a]$ for any $a > 0$
but only pointwise in $[0, \infty)$. 5

(b) State and prove the Weirstrass M -test
for uniform convergence of a series of
functions. 5

(c) State and prove the Cauchy's criterion
for uniform convergence of a sequence
of functions. 5

(d) Test the uniform convergence of—

$$(i) \sum \frac{\sin(x^2 + n^2 x)}{n(n+1)};$$

$$(ii) \sum \frac{\cos nx}{x^p} \quad 5$$

UNIT—V

9. Answer any two of the following questions :
2×2=4

(a) Find the radius of convergence of the
power series

$$1 + x + x^2 + x^3 + \dots$$

(7)

(b) Give example of a power series that is
convergent only at $x = 0$ and of another
power series that converges for all $x \in \mathbb{R}$.

(c) Given that a power series converges for
 $x = 2$, find an interval in which it is
absolutely convergent. Also, find a lower
bound for its radius of convergence.

10. Answer either [(a) and (b)] or [(c) and (d)] : 10

(a) If a power series $\sum a^n x^n$ converges for
 $x = r$, then prove that it converges
absolutely for every x with $|x| < |r|$. Give
example to show that the power series
may not be absolutely convergent at
 $x = r$. 5

(b) Determine the radius of convergence
and the exact interval of convergence of
the following power series : 3+2=5

$$(i) \sum \frac{(n+1)x^n}{(n+2)(n+3)}$$

$$(ii) \sum \frac{2^n x^n}{n!}$$

(8)

- (c) Discuss the uniform convergence of the series

$$\sum_{n=1}^{\infty} \frac{x^n}{n^\alpha}$$

for various values of α . 5

- (d) If a power series $\sum a_n x^n$ has radius of convergence R , prove that for any positive integer m the series $\sum a_n x^{mn}$ has radius of convergence $R^{1/m}$. 5
